

Mathematical Statistics

Anna Janicka

Lecture VI, 25.03.2019

PROPERTIES OF ESTIMATORS, PART II

Plan for Today

1. Fisher information
2. Information inequality
3. Estimator efficiency
4. Asymptotic estimator properties
 - consistency
 - *asymptotic normality*
 - *asymptotic efficiency*



Fisher information

If a statistical model with obs. X_1, X_2, \dots, X_n and probability f_θ fulfills regularity conditions, i.e.:

1. Θ is an open 1-dimensional set.
2. The support of the distribution $\{x: f_\theta(x) > 0\}$ does not depend on θ .
3. The derivative $\frac{df_\theta}{d\theta}$ exists.

we can define **Fisher information**

(Information) for sample X_1, X_2, \dots, X_n :

$$I_n(\theta) = E_\theta \left(\frac{d}{d\theta} \ln f_\theta(X_1, X_2, \dots, X_n) \right)^2$$



Fisher information – what does it mean?

- It is a measure of how much a sample of size n can tell us about the value of the unknown parameter θ (on average).
- If the density around θ is flat, then information from a single observation or a small sample will not allow to differentiate among possible values of θ . If the density around θ is steep, the sample contributes a lot of info leading to θ identification.



Fisher Information – cont.

Some formulae:

□ if the distribution is continuous

$$I_n(\theta) = \int_X \left(\frac{df_\theta(x)}{d\theta} \right)^2 f_\theta(x) dx$$

□ if the distribution is discrete

$$I_n(\theta) = \sum_{x \in X} \left(\frac{dP_\theta(x)}{d\theta} \right)^2 P_\theta(x)$$

□ if f_θ is twice differentiable

$$I_n(\theta) = -E_\theta \left(\frac{d^2}{d\theta^2} \ln f_\theta(X_1, X_2, \dots, X_n) \right)$$



Fisher information – cont. (2)

- If the sample consists of independent random variables from the same distribution, then

$$I_n(\theta) = nI_1(\theta)$$

where $I_1(\theta)$ is Fisher information for a single observation



Fisher Information – examples

- Exponential distribution $\exp(\lambda)$

$$I_1(\lambda) = \dots = \frac{1}{\lambda^2}$$

- Poisson distribution $\text{Poiss}(\theta)$

$$I_1(\theta) = \dots = \frac{1}{\theta}$$



Information Inequality (Cramér-Rao)

Let $X=(X_1, X_2, \dots, X_n)$ be observations from a joint distribution with density $f_\theta(x)$, where $\theta \in \Theta \subseteq \mathbb{R}$. If:

- $T(X)$ is a statistic with a finite expected value, and $E_\theta T(X)=g(\theta)$
- Fisher information is well defined, $I_n(\theta) \in (0, \infty)$
- All densities f_θ have the same support
- The order of differentiating ($d/d\theta$) and integrating $\int \dots dx$ may be reversed.

Then, for any θ :

$$\text{Var}_\theta T(X) \geq \frac{(g'(\theta))^2}{I_n(\theta)}$$


Information inequality – implications

- The MSE of an unbiased estimator (= the variance) cannot be lower than a given function of n and θ .
- If the MSE of an estimator is equal to the lower bound of the information inequality, then the estimator is MVUE.
- If $\hat{\theta}(X)$ is an unbiased estimator of θ , then

$$\text{Var}_{\theta} \hat{\theta}(X) \geq \frac{1}{I_n(\theta)}$$



Information inequality – examples

- In the Poisson model, $\hat{\theta} = \bar{X}$ is MVUE(θ) $Var_{\theta}(\bar{X}) = \frac{\theta}{n}$
- In the exponential model, \bar{X} is MVUE($1/\lambda$)

$$Var_{\lambda}(\bar{X}) = \frac{1}{n\lambda^2}$$

- The Cramér-Rao inequality is not always optimal. In the exponential model, $\hat{\lambda} = 1/\bar{X}$ is a biased estimator of λ .

$$\tilde{\lambda} = \frac{n-1}{n\bar{X}}$$

is an unbiased estimator, which is also MVUE(λ), although its variance is *higher* than the bound in the Cramér-Rao inequality.



Efficiency

The efficiency of an unbiased estimator

$\hat{g}(X)$ of $g(\theta)$ is:

$$\text{ef}(\hat{g}) = \frac{(g'(\theta))^2}{\text{Var}_{\theta}(\hat{g}) \cdot I_n(\theta)}$$

Relative efficiency of unbiased estimators

$\hat{g}_1(X)$ and $\hat{g}_2(X)$:

$$\text{ef}(\hat{g}_1, \hat{g}_2) = \frac{\text{Var}_{\theta}(\hat{g}_2)}{\text{Var}_{\theta}(\hat{g}_1)} = \frac{\text{ef}(\hat{g}_1)}{\text{ef}(\hat{g}_2)}$$



Efficiency and the information inequality

- If the information inequality holds, then for any unbiased estimator

$$ef(\hat{g}) \leq 1$$

- If $\hat{g} = \text{MVUE}(g)$, then it is possible that $ef(\hat{g}) = 1$, but it is also possible that

$$ef(\hat{g}) < 1$$

- If $ef(\hat{g}) = 1$, then the estimator is **efficient**.

Cramér-Rao efficiency



Efficiency – examples

□ In the Poisson model, $\hat{\theta} = \bar{X}$ is efficient.

□ In the exponential model, \bar{X} is an efficient estimator of $1/\lambda$.

□ In the exponential model, $\hat{\lambda} = \frac{n-1}{n\bar{X}}$

is not an efficient estimator of λ , although it is MVUE(λ).

□ In a uniform model $U(0, \theta)$, for the MLE(θ) we get $ef > 1$ (that is because the assumptions of the information inequality are not fulfilled)



Asymptotic properties of estimators

- Limit theorems describing estimator properties when $n \rightarrow \infty$
- In practice: information on how the estimators behave for large samples, *approximately*
- Problem: usually, there is no answer to the question what sample is large enough (for the approximation to be valid)



Consistency

Let $X_1, X_2, \dots, X_n, \dots$ be an IID sample (of independent random variables from the same distribution). Let $\hat{g}(X_1, X_2, \dots, X_n)$ be a sequence of estimators of the value $g(\theta)$.

\hat{g} is a **consistent** estimator, if for all $\theta \in \Theta$, for any $\varepsilon > 0$:

$$\lim_{n \rightarrow \infty} P_{\theta}(|\hat{g}(X_1, X_2, \dots, X_n) - g(\theta)| \leq \varepsilon) = 1$$

(i.e. \hat{g} converges to $g(\theta)$ in probability)



Strong consistency

Let $X_1, X_2, \dots, X_n, \dots$ be an IID sample (of independent random variables from the same distribution). Let $\hat{g}(X_1, X_2, \dots, X_n)$ be a sequence of estimators of the value $g(\theta)$.

\hat{g} is **strong consistent**, if for any $\theta \in \Theta$:

$$P_{\theta} \left(\lim_{n \rightarrow \infty} \hat{g}(X_1, X_2, \dots, X_n) = g(\theta) \right) = 1$$

(i.e. \hat{g} converges to $g(\theta)$ almost surely)



Consistency – note

From the Glivenko-Cantelli theorem it follows that empirical CDFs converge almost surely to the theoretical CDF. Therefore, we should expect (strong) consistency from all sensible estimators.

Consistency = minimal requirement for a sensible estimator.



Consistency – how to verify?

- From the definition: for example with the use of a version of the Chebyshev inequality:

$$P(|\hat{g}(X) - g(\theta)| \geq \varepsilon) \leq \frac{E(\hat{g}(X) - g(\theta))^2}{\varepsilon^2}$$

Given that the MSE of an estimator is

$$MSE(\theta, \hat{g}) = E_{\theta}(\hat{g}(X) - g(\theta))^2$$

we get a sufficient condition for consistency:

$$\lim_{n \rightarrow \infty} MSE(\theta, \hat{g}) = 0$$

- From the LLN



Consistency – examples

- For any family of distributions with an expected value: the sample mean \bar{X}_n is a consistent estimator of the expected value $\mu(\theta) = E_{\theta}(X_1)$. Convergence from the SLLN.
- For distributions having a variance:
$$S_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \quad \text{and} \quad \hat{S}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$
are consistent estimators of the variance $\sigma^2(\theta) = \text{Var}_{\theta}(X_1)$. Convergence from the SLLN.



Consistency – examples/properties

- An estimator may be unbiased but inconsistent; eg. $T_n(X_1, X_2, \dots, X_n) = X_1$ as an estimator of $\mu(\theta) = E_\theta(X_1)$.
- An estimator may be biased but consistent; eg. the biased estimator of the variance or any unbiased consistent estimator + $1/n$.





WARSAW UNIVERSITY
Faculty of Economic Sciences