

**Probability Calculus 2019/2020**  
**Problem set 12**

1. We measure a physical value 100 times. The measurement errors are independent random variables with mean 0 and variance equal to 0.1. Using the Chebyshev-Bienaymé Inequality, find an upper bound to the probability that the absolute value of the total (aggregate) error exceeds 10.
2. Using the Bernstein Inequality, find an upper bound to the probability that upon rolling a die 300 times, we will obtain a six at least 60 times.
3. Let  $X_1, X_2, \dots$  be independent random variables with distributions

(a)  $\mathbb{P}(X_n = 0) = 1/2 = \mathbb{P}(X_n = 2)$

(b) uniform over  $[0, 1]$

Does the sequence  $(Y_n)_{n \geq 1}$ , where  $Y_n = X_1 \cdot X_2 \cdot \dots \cdot X_n$ , converge in probability? Almost surely? If yes, find the limits.

4. Let  $X_1, X_2, \dots$  be uncorrelated random variables, such that  $X_n$  has a uniform distribution over  $[-1/n, 1/n]$ . Does the sequence

$$\frac{X_1 + X_2 + \dots + X_n}{n}$$

converge in probability?

5. Let  $X_1, X_2, \dots$  be independent random variables, such that  $\mathbb{P}(X_n = n) = \mathbb{P}(X_n = -n) = \frac{1}{2}$ . Does the sequence

$$\frac{X_1 + X_2 + \dots + X_n}{n}$$

satisfy the WLLN? Converge in probability to 0?

6. Let  $X_1, X_2, \dots$  be independent random variables with exponential distribution with parameter 2. Does the sequence

$$\frac{X_1 + X_2 + \dots + X_n + 3}{2n + 31}, \quad n = 1, 2, \dots$$

converge almost surely? What is the limit?

7. We randomly and independently draw points  $A_1, A_2, \dots$  from the interval  $[0, 3]$ . For a given  $n$ , let  $S_n$  denote the number of points among  $A_1, A_2, \dots, A_n$ , which fall into the interval  $[0, 1]$ . Verify, whether  $\frac{S_n}{n} \rightarrow \frac{1}{3}$  almost surely.

**Some additional simple problems you should be able to solve on your own:**

Theory (you should know going into this class)

1. Formulate the Chebyshev inequalities and the Bernstein inequality.
2. What does it mean that a sequence of random variables converges almost surely/in probability?
3. Provide the Weak and Strong Laws of Large Numbers for the Bernoulli Scheme.

Problems (you should know how to solve after this class)

- A. A symmetric coin is tossed 100 times. Using the Bernstein inequality, assess the probability that heads will appear in at least 60% cases.
- B. Let  $X_1, X_2, \dots$  be independent random variables from an exponential distribution with parameter 2. Verify, whether the sequence

$$\frac{X_1 + X_2 + \dots + X_n + 3}{n + 31}, \quad n = 1, 2, \dots$$

converges in probability.

- C. Let  $X_1, X_2, \dots$  be uncorrelated random variables, where  $X_n$  has a distribution given by  $\mathbb{P}(X_n = -n) = \mathbb{P}(X_n = n) = 1/(2n^2)$ ,  $\mathbb{P}(X_n = 0) = 1 - 1/n^2$  for  $n \geq 1$ . Verify, whether the sequence

$$\frac{X_1 + X_2 + \dots + X_n}{n}, \quad n = 1, 2, \dots,$$

converges in probability.

- D. Let  $X_1, X_2, \dots$  be independent random variables from a uniform distribution over the interval  $[0, 1]$ . Prove that the sequence

$$\frac{X_1 + X_2 + \dots + X_n}{n}, \quad n = 1, 2, \dots,$$

converges in probability and find the limit.

- E. Let  $X_1, X_2, \dots$  be independent random variables from a geometric distribution with parameter 0.1. Verify, whether the sequence

$$\frac{X_1 + X_2 + \dots + X_n - 15}{n + 31}, \quad n = 1, 2, \dots$$

converges almost surely and if yes, find the limit.

- F. Let  $X_1, X_2, \dots$  be independent random variables from a uniform distribution over the interval  $[0, 1]$ . Prove that the sequence

$$\frac{X_1 + X_2 + \dots + X_n}{n}, \quad n = 1, 2, \dots,$$

converges almost surely and find the limit.